

Leukaemia Section

Short Communication

t(2;12)(q31;p13)

Jean-Loup Huret

Genetics, Dept Medical Information, University of Poitiers, CHU Poitiers Hospital, F-86021 Poitiers, France

Published in Atlas Database: November 2006


Online updated version: <http://AtlasGeneticsOncology.org/Anomalies/t0212q31p13ID1459.html>

DOI: 10.4267/2042/38418

This work is licensed under a Creative Commons Attribution-Non-commercial-No Derivative Works 2.0 France Licence.

© 2007 *Atlas of Genetics and Cytogenetics in Oncology and Haematology*

Identity



t(2;12)(q31;p13) G- banding – Courtesy Melanie Zenger and Claudia Haferlach.

Clinics and pathology

Disease

Non Hodgkin lymphoma (NHL) and myelodysplastic syndrome (chronic myelomonocytic leukemia (CMML))

Epidemiology

Only 2 cases to date (male patients): 1 NHL aged 30 years and 1 CMML aged 78 years

Prognosis

The CMML case died 2 years 9 months after diagnosis

Cytogenetics

Additional anomalies

del(11q) in the NHL case and +21 in the CMML case.

Genes involved and Proteins

Note: ETV6 was found involved in the NHL case.

References

Sato Y, Bohlander SK, Kobayashi H, Reshmi S, Suto Y, Davis EM, Espinosa III R, Hoopes R, Montgomery KT, Kucherlapati RS, Le Beau MM, Rowley JD,. Heterogeneity in the breakpoints in balanced rearrangements involving band 12p13 in hematologic malignancies identified by fluorescence in situ hybridization: TEL (ETV6) is involved in only one half. *Blood* 1997;90:4886-4893.

This article should be referenced as such:

Huret JL. t(2;12)(q31;p13). *Atlas Genet Cytogenet Oncol Haematol*.2007;11(2):134.